

# RADIOCARBON AGE OF LATE PLEISTOCENE GLACIGENIC SEDIMENTS IN JONIONYS SECTION OF MERKINĖ (EEMIAN) INTERGLACIAL

ALGIRDAS GAIGALAS<sup>1</sup>, ANNA PAZDUR<sup>2</sup> AND JACEK PAWLYTA <sup>2</sup>

<sup>1</sup>Department of Geology and Mineralogy, Vilnius University, Čiurlionio 21/27, LT 2009 Vilnius, Lithuania  
(e-mail: Algirdas.Gaigalas@gf.vu.lt).

<sup>2</sup>Radiocarbon Laboratory, Institute of Physics, Silesian University of Technology,  
Krzywoustego 2, PL 44-100 Gliwice, Poland  
(e-mail: Pazdur@zeus.polsl.gliwice.pl; pawlyta @polsl.gliwice.pl)

## Key words:

RADIOCARBON AGE,  
ESR AGE,  
OSL AGE,  
LATE PLEISTOCENE,  
GLACIGENIC  
SEDIMENTS,  
JONIONYS SECTION,  
STRATIGRAPHY,  
LITHUANIA

**Abstract:** The stratotype of Merkinė (Eemian) interglacial in Lithuania is the Jonionys section near Merkinė town in south Lithuania. The lacustrine sediments containing rich fossil flora in this section were formed under the conditions of Merkinė interglacial and Nemunas (Vistulian) glacial. The fossil-bearing Merkinė deposits are overlain by a probably complete sequence of Early Nemunas and partly of Middle Nemunas deposits. Rewashed till-covered organic sediments were found in the upper part of Jonionys section. The exact age of glacial sediments of this section has remained largely unknown. Peat with wood remnants under the rewashed till have been investigated in Gliwice Radiocarbon Laboratory (Gd-10825: 31,500+2300/-1800 BP and Gd-14000: >31,000 BP). Its deposition probably took place just before the maximum of the last glaciation. In Jonionys section the Merkinė interglacial, Early and Middle Nemunas nonglacial sediments were probably accumulated in the time of climatic fluctuations, but without glacigenic sedimentation. The glacigenic sediments in the upper part of Jonionys stratotype section are younger than 30,000 BP and belong to the Late Nemunas glacial maximum. Recent radiocarbon dating are important for the establishment of the standard area of the European Würm (Vistulian, Valdaian, Poozerian and Nemunas) glacial.

## 1. INTRODUCTION

The last Merkinė (Eemian) interglacial and last Nemunas (Vistulian) glacial macrocycle of the Late Pleistocene (from 130,000 to 10,000 BP) in Lithuania is probably more intensely studied, than the other subdivisions of the Pleistocene. There are three stages of the last glaciation: Early (70,000-55,000 BP), Middle (55,000-30,000 BP) and Late (30,000-10,000 BP) (Gaigalas and Melešytė, 1999). The glacial activity was different at each stage. The data already obtained in the Jonionys section confirm the absence of an ice sheet in south Lithuania during the Early and Middle Nemunas (Gaigalas, 1995; Gaigalas and Hütt, 1996). The ability to date limno-alluvial sediments and soils by methods of absolute chronology provides the opportunity to establish reliable geochronology of the Upper Pleistocene in Lithuania (Gaigalas, 2000).

Merkinė interglacial deposits are presented by lake-bog deposits (gyttja, peat, clay, mud, loam and sand). The thickness of these deposits is variable. In some sections

they are 2-3 m thick and in other they reach more than 20 meters. The Merkinė interglacial deposits are most widely spread. They were found in different areas of Lithuania except the west part near the seaside. Most sections have been examined in the south-eastern Lithuania between Merkinė and Druskininkai.

The region between Merkinė and Druskininkai is considered to be a stratotypical region and the deposits of Jonionys are assigned to be stratotypical ones (Fig. 1). The sections characteristic of the Merkinė (Eemian) interglacial are most frequently found in the vicinity of the town of Merkinė. The name of the Merkinė interglacial was taken from the name of this town (Kondratienė, 1965; Vaitiekūnas, 1968). The Merkinė interglacial stratotype deposits are located in the Jonionys-Maksimionys site (Fig. 1). At present more than 30 sections of the Merkinė interglacial were studied in Lithuania. Parastratotype sections are known from Netiesos, Kibyšiai, Druskininkai (Pušynas and park), Ratnyčia, Liškiava, Giraitiškės, Smalininkai, Kmitos, Puponyš-674, Medininkai (bore-

holes 2, 117, 3), Mickūnai-Gaidūnai (boreholes 5, 7, 9), Bezdonyš – 296 and 78, Arvydai – 79, Skersabaliai – 110, Kurkčiai, Bukiškės-25 and 80, Gervelė-330, Gaurė and others.

## 2. SITE DESCRIPTION OF MERKINĖ INTERGLACIAL

The formation age of the Jonionys and many other sections comprises not only the Merkinė interglacial, but also Medininkai Glaciation and the Nemunas Glaciation.

The studied Jonionys section is an open natural exposure located on the left bank of the Nemunas River about 3 km west of Merkinė (south Lithuania). It reveals the socle of the second above flood plain terrace of the Nemunas River. The geological section (Fig. 2), from the bottom (above mean water level of the Nemunas River) to top, is the following:

- 1 – gIIImd – till (morainic loam) of Middle Pleistocene, Medininkai Glaciation, of grayish brown color, 0.4 m;
- 2 – I gIIImd-IIIImr – sandy silt of the end of the Medininkai Glaciation and beginning of the Merkinė interglacial ( $M_1$  – *Pinus sylvestris*, *Picea obovata* and *Betula*), 0.2 m;
- 3 – IIIImr – clayey-silty sand with gravel, of dark gray color, with small shell of mollusks and detritus ( $M_2$ – $M_{3a}$  – *Quercus petraea*, *Quercus pubescens*, *Pinus sylvestris* *Ulmus glabra*, *Ul. levis*, *Caulina flexilis* and *Scirpus lacustris*), 0.4 m;
- 4 – IIIImr – sand of yellowish gray color, middle-fine grain size, humic, with small shell of mollusks and detritus ( $M_{3b}$  – *Tilia tomentosa*, *T. platyphyllos*, *T. cordata*, *Vitis*, *Hedrea*, *Acer compestre*, *A. platanoides*, *Alisma plantago aquatica* and *Ceratophyllum demersum*), 0.5 m;

- 5 – IIIImr – silty sand of grayish brown color, fine grain size, humic, with wood remains ( $M_{3c}$  – *Carpinus betulus*, *Corylus avellana*, *Tilia tomentosa*, *T. platyphyllos*, *Caulina flexilis*, *Lemna trisulca*, *Osmunda cinnamomea* and *O. Regalis* and *Ligustrum*, *Vitis*), 0.4 m;
- 6 – IIIImr – sandy gyttja of grayish brown color, with small wood detritus ( $M_4$  – *Picea obovata*, *Carpinus betulus*, *Swida sanguinea* and *Osmunda cinnamomea*), 0.2 m;
- 7 – IIIImr-IIIInm<sub>1</sub> – sandy silty clay of grayish blue color (lower part, about 0.1 m – (Merkinė interglacial),  $M_5$  – *Pinus sylvestris* (upper part, about 0.3 m – lower Nemunas glacial), Nm<sub>1a</sub> – *Pinus sylvestris*, *Alnus glutinosa*, *Acea obovata*, *Selaginella selaginoides*, *Sparganium simplex*, *Chenopodium album* and *Menyanthes trifoliata*; with rare shells of mollusks and small insertion of wood remains, 0.4 m. On the top of this sequence of Merkinė deposits of Jonionys (Fig. 2) there is the following succession of Early Nemunas sediments:
- 8 – IIIInm<sub>1</sub> – silt with lenses of sand, of dark gray color, humic, with rare carbonization remains of plants ( $J_1$  – *Juniperus communis*), 0.1 m;
- 9 – IIIInm<sub>1</sub> – clayey sand silt, of gray color with brown tint, humic, horizontally stratified ( $J_1$  – *Juniperus communis* and *Potamogeton perfoliata*), 0.15 m;
- 10 – IIIInm<sub>1</sub> – silty sand, of gray color with brown tint, humic ( $J_1$  – *Betula alba*, *Alnus glutinosa*, *Pinus sylvestris*, *Potamogeton filiformis*, *Scirpus lacustris* and *Lycopodium clavatum*), 0.1 m;
- 11 – IIIInm<sub>1</sub> – silty clay, of gray color, humic, massive structure ( $J_1$  – *Betula alba*, *Alnus glutinosa*, *Pinus sylvestris*, *Potamogeton filiformis*, *Scirpus lacustris* and *Lycopodium clavatum*), 0.15 m;




Fig. 1. Location of the Jonionys section near the Merkinė town in stratotype region Druskininkai-Merkinė (stratotype palaeobasin of Jonionys-Maksimonys) with geochronological limits of Grūda Stadial (G), Žiogeliai Phasial (Ž), Baltija Stadial (B), South-Lithuanian (SL), Middle-Lithuanian (ML) and North-Lithuanian (NL) Phasials.

- 12 – IIIInm<sub>1</sub> – ferruginous sand, of gray color with brown tint, middle-fine grain size, massive structure, (Nm<sub>1b</sub> – *Najas marina* and *Urtica dioica*), 0.2 m;
- 13 – IIIInm<sub>1</sub> – sand of brown color, fine-middle grain size, horizontally stratified, with lamina of silt (Nm<sub>1b</sub> – *Selaginella selaginoides*, *Hippuris vulgaris*, *Lycopodium alpinum*, *Botrychium boreale* and *Betula nana*), 0.15 m;
- 14 – IIIInm<sub>1</sub> – sandy silty clay, of gray color with brown tint, massive structure (Nm<sub>1b</sub> – *Alnus glutinosa*, *Thalictrum*, *Betula nana*, *Lycopodium alpinum*, *Selaginella selaginoides*, *Botrychium boreale*, *Salix*, *Alnaster*, *Cyperaceae*, *Gramineae* and *Chenopodiaceae*), 0.15 m;
- 15 – IIIInm<sub>1</sub> – sand of gray color with brown tint, with laminae of silt and ripples of flow, weakly humic, with rare detritus, in lower part shells of *Anadonta* (Nm<sub>1b</sub> – J<sub>2</sub> – *Betula alba* and *Pinus sylvestris*; fauna of rodents: *Apodemus sp.*, *Arvicola cf. terrestris* L., *Lemmus sibiricus* Kerr., *Microtus sp.*, *M. agrestis* L. and *M. arvalis* Pall.), 0.8 m;
- 16 – IIIInm<sub>2</sub> – sand of lightly gray color, in the upper part – yellowish, with laminae of darkly brown gyttja (R<sub>1</sub> – *Betula alba*, *B. nana*, *B. pubescens*, *Polygonum sp.*, *Pinus sylvestris*, *Quercus sp.*, *Cyperaceae*, *Gramineae*, *Lycopodium clavatum* and *L. annolinum*; fauna of rodents: *Sorex sp.*, *Arvicola aff. terrestris* L., *Microtus oeconomus* Pall., *M. sp.*, *Pitymys of subterraneus* Sel.-Long. and *Alces sp.*), 0.4 m;
- 17 – IIIInm<sub>2</sub> – clayey silty sand of dark gray color, weakly humic, the upper contact is eroded (R<sub>2</sub> – *Betula nana*; fauna of rodents: *Polygonum lapathifolium*), 0.1 m;
- 18 – IIIInm<sub>3</sub> – sand of yellow color, mixed grain size, with allochthonous pieces of gyttja, the upper contact is eroded, 0.1 m;

- 19 – gIIIInm<sub>3</sub> – rewashed till sand-gravel-pebble sediments with simple boulders, fragments of sedimentary rocks, prevail well-rounded, 0.25 m;
- 20 – silty sand of lightly yellowish gray color, fine grain size, 2.0 m.

The deposits of lake and bog origin have formed since the beginning of the interglacial up to the last glaciation in Jonionys site (Fig. 3). It has been established that the sediments in this location were formed under the conditions of Merkinė interglacial and Nemunas glacial (Kondratienė, 1996).

The results of the investigation at the Jonionys site illustrate the presence of three thermometers younger than the Eemian interglacial. The first two (correlated with Brörup and Odderade) are, according to recent data, rather widely represented in Lithuania and recently have been identified at several other sites (Satkūnas, Grigienė and Robertsson, 1998). The Jonionys-3 thermometer is separated from the preceding one by a cryomer layer formed under nonglacial condition. The Jonionys-3 interstadial corresponds to the Oerel interstadial. The cryomer Nemunas 2a marks the beginning of the isotope stage 4. This stage by some researchers is interpreted as representing a glacial advance. According to the data from southern and eastern Lithuania, no till beds that can be attributed to the beginning of the Middle Nemunas have been found.

The exact age of glacial sediments of this section has remained largely unknown. The Merkinė interglacial site at the stratotype locality of Jonionys is represented by lacustrine deposits containing a rich fossil flora. The lower boundary of Nemunas deposits can be defined on palaeobotanical criteria. The Nemunas ice-free deposits in the Jonionys section overlie the sediments of the




Fig. 2. Stratigraphic sequence of the Merkinė and Nemunas deposits exposed at Jonionys with OSL (in ka), ESR and <sup>14</sup>C dates: 1 – crystalline rocks, 2 – sandstones and siltstones, 3 – dolostones, 4 – Ordovician and Silurian limestones, 5 – other limestones, 6 – Mesozoic marls. Structural diagrams of pebble long axes orientation in tills.



area north of Grodno and Raduny as well as north of Minsk along the Ostrovets-Dokshitsy-Orsha line (Matveyev, 1995). Maximum range of the ice sheet during the Vistulian (Nemunas and Poozerian) Glaciation has been also reconstructed by T. Krzywicki (2001). During the Vistulian Glaciation the glacier advanced to the area of eastern Mazury, Suwałki and Grodno district during the Świecie stadial (*ca* 55,000-67,000 BP) and during the main stadial (*ca* 24,000-12,000 BP). The ice sheet limit during the Świecie (Varduva stadial and Ozierska phase). The phasial in this area was more extensive about 30 km further to the south of the limit of the main stadial (Krzywicki, 2001).

#### 4. <sup>14</sup>C AGE OF MAXIMUM NEMUNAS GLACIATION IN LITHUANIA

Evidently, the ice sheet did not penetrate into the southeastern Lithuania region during the Early and Middle Nemunas (Valdaian, Vistulian and Weichselian) interval. Ice-free interval at the Early and Middle of the Vistulian (Weichselian and Valdaian) were identified in many sites of Middle and East Europe, where the last glacial maximum took place between 25,000 and 15,000 BP. In Lithuania lithological research led to the recognition of two independent Late Pleistocene Grūda and Baltija till horizons (Gaigalas, 1998, 2001; Gaigalas and Melešytė, 1999, 2001), which were correlated with the Brandenburgian (Leszno) and Pomeranian (Pomorze) glacial stages. This subdivision was confirmed by recent <sup>14</sup>C dating. The southeastern part of Lithuania was not covered by Late Nemunas glacier. The deposits of the Nemunas Glaciation maximum are not of glacial origin and are presented by periglacial sediments – glaciolimnic silts and clay (25,000-15,000 BP). The data (Gaigalas and Satkūnas, 1996) obtained so far from the study of the Medininkai section (borehole 117A) confirm the absence of an ice sheet in this region during the Early, Middle and Late Nemunas. The glaciolimnic silt and clay covered lacustrine and peat sediments with organics were probably accumulated during the interval from 29,800 ± 1200 to 23,770 ± 630. Their deposition probably took place during and just before the last glacial advance. The glaciolimnic silt and clay in Medininkai highland are simultaneous to maximum of the last ice cover (23,000-15,000 BP). Here and there between the upper glaciolimnic silt and clay sediments and the Merkinė interglacial sediments, layers of Early and Middle Nemunas deposits occur. In this section the Merkinė interglacial, Early and Middle Nemunas nonglacial sediments were probably accumulated during the interval from 114,000 to 23,000 BP, a period with climatic fluctuations, but without glacial sedimentation. Radiocarbon dating have confirmed the conclusions received by palynological research (Kondratienė, 1996). There were more palynological data indicating the presence of organic sediments of the Merkinė interglacial, lying on the surface of relief, not covered by the glacial sediments (tills) of the Late Pleistocene.

The south-eastern part of Lithuania was not covered by continental ice of either the Grūda or Baltija stadials

of the Late Nemunas glacial. The petrographical composition of till of the Grūda stadial of Nemunas glacial in Lithuania indicates a Central Swedish source for erratic boulders and clastic material (Gaigalas, 1995). The Grūda till was enriched with local Mesozoic sedimentary rocks. Apart from Mesozoic marls clasts, this till is rich in crystalline rocks from Central Sweden, the Åland Island and the Baltic Sea floor. The Grūda ice sheet moved therefore from the north-west to the south-east. During the retreat a minor readvance of glacier, the Žiogeliai Phasial, occurred. The Baltija till deposited by the stadial glacier advance notably contains increased frequency of dolostones derived from Devonian rocks of the east Baltic region. The ice of the Baltija stadial transgressed the area of Lithuania in three distinct lobes: west Lithuanian, central Lithuanian and east Lithuanian. The glacial deposits of each of the three lobes differ in the petrographic composition of the erratics of crystalline rocks and clasts of sedimentary rocks. However, the main mass of the ice moved from the north to the south over east Baltic Palaeozoic rocks. The till of the Baltija stadial has specific association of Palaeozoic sedimentary rocks from east Baltic region and crystalline rocks from south Finland. Retreating and periodically recessing, the glacier of the Baltija stadial has left its phasial tills, which were distributed by tracks of East-Lithuanian (about 16,000 BP), South-Lithuanian (*ca* 15,000 BP), Middle-Lithuanian (*ca* 14,000 BP), and North-Lithuanian (*ca* 13,000 BP) phasials on the surface of Lithuania (Fig. 1). The East-Lithuanian phasial corresponds to the maximum of the Baltija stadial (Fig. 1). In the zone of the south Lithuanian recessional phase, the three oscillational branches of end moraines have been found. Immediately after the each retreat of the ice sheet, vast areas of Lithuania were covered with big ice-dammed lakes in interphasials. The phasial tills are spread locally, in recession zones of phasial glaciers, so their full lithostratigraphical sequence may be revealed only by carrying out successive investigation and correlation of Baltija tills on the whole area of Lithuania.

#### 5. CONCLUSIONS

The radiocarbon dating provides a more accurate chronology of the Late Nemunas (Vistulian) Glaciation. In Jonionys section reworked till with boulders of the last ice cover of the Late Nemunas Glaciation is younger than 30,000 BP. The main Late Nemunas glacier covered the Baltic countries and reached south-eastern Lithuania about 22,000-20,000 BP. In Lithuania lithological research led to the recognition of two independent Late Pleistocene Grūda (before 22,000-18,000 BP and Baltija (16,000-13,000 BP) till horizons, which were correlated with the Brandenburgian and/or Leszno and Pomeranian and/or Pomorze glacial stages. The covering till strata at the vicinity of Jonionys belongs to maximum of the Late Nemunas Glaciation. The recent radiocarbon dating is important for the establishment of the studied area as a standard for the European Würm (Vistulian, Weichselian, Valdaian, Poozerian, Nemunas) Glacial.

## REFERENCES

- Arslanov Ch.A., 1987:** *Radiocarbon: geochemistry and geochronology* (in Russian). Leningrad University: 298.
- Gaigalas A., 1995:** Glacial history of Lithuania. In: *Glacial deposits in north-east Europe*: 127-135.
- Gaigalas A., 1998:** The evolution of geological environment of the castles of Vilnius. *PACT* 54(I.10): 111-130.
- Gaigalas A., 2000:** Correlation of  $^{14}\text{C}$  and OSL dating of Late Pleistocene deposits in Lithuania. *Geochronometria* 19: 7-12.
- Gaigalas A., 2001:** Stratigraphy and geochronology of Upper (Late) Pleistocene (in Lithuanian). In: *Stone age in South Lithuania (according to geological, palaeogeographical and archaeological data)*. Vilnius: 7-24.
- Gaigalas A. and Hütt G., 1996:** OSL dating of the Merkinė (Eem) Interglacial (in Jonionys) and Nemunas Glaciation (Rokai section) in Lithuania. *PACT* 50: 59-69.
- Gaigalas A. and Melešytė M., 1999:** Stratigraphy and geochronology of Last Glacial in Lithuania. In: *The Fourth Baltic Stratigraphical Conference. Problems and methods of modern regional stratigraphy. Abstracts*. Riga: 22-24.
- Gaigalas A. and Melešytė M., 2001:** Spreading and composition of glacial deposits of the Nemunas Glaciation (in Lithuanian). In: *Stone age in South Lithuania (according to geological, palaeogeographical and archaeological data)*. Vilnius: 46-54.
- Gaigalas A., Molodkov A. and Melešytė M., 1994:** The first ESR dating results of Butėnai (Likhvin) and Merkinė (Mikulino) Interglacial deposits in Lithuania. In: *Conference on Geochronology and Dendrochronology of Old Town's and Radiocarbon Dating of Archaeological Findings Lithuania. Abstracts and Papers*. Vilnius: 15.
- Gaigalas A. and Satkūnas J., 1996:** Application of radiocarbon dating for mapping of the limits of the Last Glaciation (Nemunas, Weichselian) in South-East Lithuania. *Geologija* 19. Vilnius: 26-30.
- Kondratienė O., 1965:** Stratigraphisch ausgeführte Einteilung pleistozäner Ablagerungen Südostlitauens auf Grund palinologischer Untersuchungsergebnisse (in Russian). In: *Stratigraphie Quartärer Ablagerungen Südostlitauens und Antropogäne Paläogeographie. Arbeiten II Bd.* Mintis, Vilnius: 189-261.
- Kondratienė O., 1996:** *The Quaternary stratigraphy and palaeogeography of Lithuania based on palaeobotanic studies* (in Russian). Vilnius: 213.
- Krzywicki T., 2001:** Maximum ice sheet limit of the Vistulian Glaciation in north-eastern Poland and neighbouring areas. *Field symposium on Quaternary Geology in Lithuania. Abstract volume*. Vilnius: 41-42.
- Lindner L. and Marks L., 1995:** Outline of palaeogeomorphology of the Polish territory during the Scandinavian glaciations. *Przegląd Geologiczny* 43 (7): 591-594 (in Polish).
- Lindner L. and Yelovicheva Y.K., 1998:** First tentative correlation scheme of glacial and interglacial units in the Pleistocene of Poland and Belarus. *Quaternary Studies in Poland*, 15: 27-35.
- Matveyev A.V., 1995:** Glacial history of Belarus. In: *Glacial deposits in north-east Europe*: 267-276.
- Mojski J.E., 1959:** On the stratigraphy of loess near Hrubieszów (Lublin Upland). *Przegląd Geologiczny* 7(10): 447-450 (in Polish)
- Mojski J.E., 1985:** *Geology of Poland*. Vol. I: *Stratigraphy*, Part 3 b: *Cainozoic. Quaternary* (in Polish). Warszawa: 244.
- Satkūnas J., Grigienė A. and Robertsson A.M., 1998:** An Eemian – Middle Weichselian sequence from the Jonionys site, Southern Lithuania. *Geologija* 25, Vilnius: 82-91.
- Vaitiekūnas P., 1968:** Stratigraphical problems of Neopleistocene in Lithuania (in Polish). *Kwartalnik Geologiczny* 12(3): 646-664
- Wysota W., 1999:** Ice sheet maximum limit of the Vistulian Glaciation in the mid-eastern Chełmno-Dobrzyń Lakeland, northern Poland. *Geological Quarterly* 43(2):189-202.
- Wysota W., 2001:** Chronology of the Vistulian glacial events in the Lower Vistula region. In: *Book of Abstracts, 7<sup>th</sup> International Conference "Methods of Absolute Chronology", 23-26th April, Ustroń, Poland*: 63.

The Eemian (also called the last interglacial, Sangamonian Stage, Ipswichian, Mikulin, Kaydaky, penultimate, Valdivia or Riss-WÄ¼arm) was the interglacial period which began about 130,000 years ago at the end of the Penultimate Glacial Period and ended about 115,000 years ago at the beginning of the Last Glacial Period. It corresponds to Marine Isotope Stage 5e. Although sometimes referred to as the "last interglacial" (in the "most recent previous" sense of "last"), it was the second-to-latest. Sediment ages were determined by combining radio-carbon AMS analyses with OSL dating on quartz and feldspar in order to cover the entire expected age range and to obtain ages of sediment with only a few plant re-mains and low organic carbon contents. Table 1 and Fig. 28.1 show the dating of plant remains found in the upper section of Unit II between 44.7 and 44.8 m b.s.l. (composite ice-wedge horizon) gave a radiocarbon age of 43.815.5/±3.2 ka BP. Eemian Interglacial (Unit I). The lowermost and oldest Unit I was encountered only in core C2 and is probably of marine origin, based on high pore-water salinities, its relatively fine grain-size composition, fossil diatom records and pore-water hydrochemistry. Eemian/Kazantzevo sediments in the Ozernaya River sections on southern October Revolution Island (MIS 5e; marine unit M-III, Fig. 28.2) form a regression sequence, initiated by off-shore glaciomarine marine sediments capped by shoreface to foreshore sands and gravels, in turn capped by till (till unit T-IV) of obvious Weichselian age (Müller et al., 2007). Both marine and continental Eemian interglacial deposits have been found in Estonia. However, palynological and diatom investigations have confirmed an Eemian age of these deposits and their correlation to other Eemian sections (Cheremisinova, 1961; Raukas, 1978; Liivrand, 1984, 1991; Kondratiene, 1993; Donner, 1995).