

Corticosteroid-induced mania: Prepare for the unpredictable

Head off this common psychiatric side effect

Michael A. Cerullo, MD

Clinical assistant professor of psychiatry
Division of bipolar disorders research
University of Cincinnati College of Medicine
Cincinnati, OH

Can corticosteroids “unlock” hidden potential for mania, or are steroid-induced mood symptoms a temporary reaction? And when these mood symptoms occur, what is the best way to treat them?

Psychiatric symptoms develop in 5% to 18% of patients treated with corticosteroids. These effects—most often mania or depression—emerge within days to weeks of starting steroids. To help you head off manic and mixed mood symptoms, this paper examines how to:

- treat steroid-induced mania or mixed bipolar symptoms
- reduce the risk of a mood episode in patients who require sustained corticosteroid therapy.

‘STERIOD PSYCHOSIS’

Jane Pauley, NBC’s *Today Show* broadcaster, described in her autobiography how hypomania developed within weeks after she started cortico-



© 2006 Roy Scott



steroids for idiopathic urticaria edema: "I was so energized that I didn't just walk down the hall, I felt like I was motoring down the hall. I was suddenly the equal of my high-energy friends who move fast and talk fast and loud. I told everyone that I could understand why men felt like they could run the world, because I felt like that. This was a new me, and I liked her!"¹

Pauley's hypomania led to a manic episode and eventually to depression. She was started on antidepressants, which triggered another manic episode. Pauley—who had no history of bipolar disorder—spent 3 weeks in a New York psychiatric hospital.¹

Diagnostic symptoms. Corticosteroids' psychiatric effects—cognitive, mood, anxiety, and psychotic symptoms—were first described as "steroid psychosis." Psychosis can occur, but mood symptoms are more common:

- Among 122 patients, 40% experienced depression, followed by mania (28%), psychosis (14%), delirium (10%), and mixed mood episodes (8%).²
- Among 130 patients, mania was most prevalent (35%), followed by depression (28%), mixed mood episodes (12%), delirium (13%), and psychosis (11%).³
- Corticosteroids caused 54% of organic mania cases on a hospital psychiatric consult service.⁴
- In a prospective study of 50 patients treated with corticosteroids, 13 developed hypomania and 5 developed depression.⁵

Steroid-induced symptoms emerge from 3 to 4 days to a median of 11 days after a patient starts corticosteroid therapy. After steroids are discontinued, depressive symptoms persist approximately 4 weeks, mania 3 weeks, and delirium a few days. Approximately one-half of patients with steroid psychosis improve in 4 days and one-half within 2 weeks.^{2,6}

Risk of psychiatric symptoms was 1.3% with dosages <40 mg and 18.4% with dosages >80 mg

WHO IS AT RISK?

Corticosteroids include the steroids produced in the adrenal gland (such as corticosterone) and their synthetic—and often more potent—analogs (such as prednisone).⁷ Because of their glucocorticoid, immunosuppressant, mineralocorticoid, and anti-inflammatory properties, steroids are used as replacement therapy and to treat a wide variety of illnesses (*Table 1*).

Age and gender. Patient age appears unrelated to development of psychiatric symptoms after corticosteroid use.² One study suggested women are twice as likely as men to develop psychiatric symptoms (77 versus 38 cases in 115 patients),³ but many illnesses that require corticosteroid treatment occur more frequently in women. Other researchers found a slight female predominance (58% versus 42% of cases) when they excluded patients with systemic lupus erythematosus and rheumatoid arthritis, which are more common in women than in men.²

Dosage. Higher corticosteroid dosages increase the risk of psychiatric symptoms. In patients taking prednisone, the Boston Collaborative Drug Surveillance Project⁸ found the incidence of psychiatric side effects to be:

- 1.3% in patients taking <40 mg
- 4.6% in those taking 41 to 80 mg
- 18.4% in those taking >80 mg.

Psychiatric history. Past psychiatric illness does not seem to be a risk factor for psychiatric side effects of corticosteroids,⁹ although patients with a history of posttraumatic stress disorder are more likely to suffer depression while taking corticosteroids.¹⁰

Corticosteroid exposure. Patients who did not experience psychiatric side effects with corticosteroids in the past appear not to be protected if corticosteroids are used again. One report examined 17 cases of steroid-induced psychiatric illness in

continued on page 47

continued from page 44

patients with previous exposure to corticosteroid therapy. Six patients had previous psychiatric side effects while taking corticosteroids, and 11 did not.²

BIPOLAR TRIGGER?

Do corticosteroids' acute psychiatric side effects have long-term sequelae? Longitudinal evidence is scarce, but a few reports suggest corticosteroids could play a role in the onset of primary bipolar I disorder:

- A 28-year-old woman with no known mood symptoms before a short course of prednisone experienced six episodes of mania and depression when not taking corticosteroids during the subsequent 18 months.¹¹

- Among 16 patients with first-onset mood symptoms after corticosteroid use, a retrospective chart review found 7 had recurrent manic and depressive symptoms unrelated to additional corticosteroid use.¹²

Although intriguing, these case reports are inconclusive. Because bipolar type I incidence in the general population is 1.5%,¹³ many persons with bipolar disorder undergo corticosteroid treatment. Nevertheless, these results—especially from the retrospective review¹²—suggest that corticosteroid use may contribute to the onset of bipolar I illness.

SYMPTOMATIC TREATMENT

Corticosteroid-induced side effects are usually managed by tapering off the steroids and treating the psychiatric symptoms.^{2,3} Simply tapering off the steroids—without additional treatments—led

to recovery in 33 of 36 patients.² Stopping corticosteroids is not always possible or desirable, however, especially in many medically complicated cases seen by psychiatric consult services.

In a recent case, I was asked to see a man, age 69, on the oncology service who was receiving corticosteroids every 2 weeks as part of his chemotherapy. The patient was admitted to the hospital for acute mental status changes 2 days after his last corticosteroid dose. He had pressured speech, grandiosity, and had not slept in 2 days. We started risperidone, 1 mg bid, and most of his manic symptoms resolved within 2 days. His chemotherapy was continued without corticosteroids. If this had not been possible, I would have recommended continuing risperidone prophylactically.

Table 1

Medical conditions for which corticosteroids are commonly used

Disorder	Indications for corticosteroids
Acute adrenal insufficiency	Acute; replacement therapy
Addison's disease	Chronic; replacement therapy
Asthma	Acute and chronic; anti-inflammatory
Inflammatory bowel disease	Acute; anti-inflammatory
Multiple sclerosis	Acute; exacerbations, immunosuppressant
Organ transplant	Chronic; immunosuppressant
Rheumatoid arthritis	Chronic; anti-inflammatory
Systemic lupus erythematosus	Acute; severe exacerbation, immunosuppressant (high doses are used)

continued



Table 2

Mood stabilizers with evidence of benefit in treating corticosteroid-induced mania

Indication	Medication	Dosage/blood level	Evidence
Preventing psychiatric effects in patients requiring long-term corticosteroids	Lithium	0.8 to 1.2 mEq/L	Prospective trial (27 with multiple sclerosis) ²⁴
Preventing recurrence of manic symptoms in patients requiring additional steroid pulses	Carbamazepine	600 mg qd (to therapeutic range of 8 to 12 µg/mL)*	Case report ¹⁶
	Gabapentin	300 mg tid	Case report ²⁶
Treating steroid-induced manic symptoms	Olanzapine	Initially 2.5 mg/d, titrated to 20 mg/d	Open-label trial (12 patients) ¹⁴
	Lithium	0.7 mEq/L	Case report ¹⁵
	Quetiapine	25 mg qhs and 12.5 mg bid prn	Case report ¹⁷
	Carbamazepine	600 mg qd (to therapeutic range of 8 to 12 µg/mL)*	Case reports ^{12,16}
Treating steroid-induced depressive symptoms	Haloperidol	2 to 20 mg/d*	Case reports ^{12,16}
	Fluoxetine	20 mg/d	Case report ¹⁸
	Amitriptyline	30 mg/d (usual effective range is 50 to 300 mg/d)*	Case report ¹²
	Lamotrigine	Up to 400 mg/d	Case report ¹⁹
Treating steroid-induced psychotic symptoms	Lithium	0.1 to 0.8 mEq/L	Case reports ^{20,21}
	Haloperidol	5 mg IV on day 1, then 2 mg po bid	Case report ²²
	Risperidone	1.5 mg/d	Case report ²³

* Dosage not included in published report; recommendation based on experience or anecdotal information

No double-blind, placebo-controlled studies have examined prevention or treatment of steroid-induced mania or other psychiatric symptoms. Uncontrolled trials and case reports suggest benefit from some symptomatic and preventive treatments (Table 2).

Treating manic and mixed mood symptoms. Twelve outpatients with manic or mixed symptoms from corticosteroid use received olanzapine in a 5-week, open-label trial. Flexible dosing started at 2.5 mg/d

and was increased as needed (maximum 20 mg/d). One patient dropped out for lack of efficacy. For the others, manic and mixed symptoms improved significantly, as indicated by scores on the Young Mania Rating Scale, Hamilton Rating Scale for Depression, and Brief Psychotic Rating Scale.¹⁴ Patient weight, blood glucose, and involuntary movements did not change significantly.

Evidence from case reports indicates that lithium,¹⁵ carbamazepine,^{12,16} haloperidol,^{12,16} or

quetiapine¹⁷ also can successfully treat steroid-induced manic symptoms.

Treating other psychiatric symptoms. Case reports support electroconvulsive therapy,^{2,15} fluoxetine,¹⁸ amitriptyline,¹² lamotrigine,¹⁹ or lithium^{20,21} for steroid-induced depression, and haloperidol²² or risperidone²³ for steroid-induced psychosis.

In four cases,⁶ tricyclic antidepressants appeared to worsen corticosteroids' psychiatric side effects. These case patients might have had steroid-induced delirium instead of mood disorders or psychosis, however, and the tricyclics' anticholinergic effects could have worsened the delirium.⁹

PREVENTING STEROID-INDUCED SYMPTOMS

Although clear guidelines on when to start preventive treatments do not exist, potential candidates for pretreatment with lithium or other agents include patients who:

- have developed psychiatric symptoms multiple times after repeated corticosteroid use
- are at high risk if psychiatric side effects occur.

Lithium. Prophylactic lithium was given to 27 patients with multiple sclerosis and taking corticosteroids for acute exacerbations. None developed psychiatric symptoms.²⁴ At the same clinic, 6 of 44 patients with multiple sclerosis or retrobulbar neuritis developed psychiatric side effects after using corticosteroids without lithium.

Be cautious when using prophylactic lithium because some conditions treated with corticosteroids—such as systemic lupus erythematosus—can impair renal function.²⁰ Corticosteroids also can affect sodium balance and increase the risk of lithium intoxication.²⁵

Check renal function before and during lithium titration, and initiate corticosteroid therapy when lithium is at effective blood levels (0.8 to 1.2 mEq/L). Monitor lithium levels and renal function frequently during steroid treatment.

Other mood stabilizers. Two case reports describe patients who repeatedly developed manic symptoms after multiple corticosteroid doses. Carbamazepine, 600 mg qd,¹⁶ and gabapentin, 300 mg tid,²⁶ prevented manic symptoms after additional corticosteroid pulses.

References

1. Pauley J. *Skywriting: a life out of the blue*. New York: Random House; 2004.
2. Lewis DA, Smith RE. Steroid-induced psychiatric syndromes. A report of 14 cases and a review of the literature. *J Affect Disord* 1983;5:319-32.
3. Sirois F. Steroid psychosis: a review *Gen Hosp Psychiatry* 2003; 25:27-33.
4. Rundell JR, Wise MG. Causes of organic mood disorder. *J Neuropsychiatry Clin Neurosci* 1989;1:398-400.
5. Naber D, Sand P, Heigl B. Psychopathological and neuropsychological effects of 8-days' corticosteroid treatment. A prospective study. *Psychoneuroendocrinology* 1996;21:25-31.
6. Hall R, Popkin M, Stickney S, Gardner E. Presentation of the steroid psychoses. *J Nerv Ment Dis* 1979;167:229-36.
7. Schimmer B, Parker K. Adenohypophyseal hormones and their hypothalamic releasing factors. In: Hardman J, Limbird L, Gilman A (eds). *Goodman and Gilman's the pharmacological basis of therapeutics, 9th ed*. New York: McGraw-Hill;1996:1459-86.
8. The Boston Collaborative Drug Surveillance Program. Acute adverse reactions to prednisone in relation to dosage. *Clin Pharmacol Ther* 1972;13:694-8.
9. Patten SB, Neutel CI. Corticosteroid-induced adverse psychiatric effects: incidence, diagnosis and management. *Drug Saf* 2000; 22:111-22.
10. Brown ES, Suppes T, Khan DA, Carmody TJ 3rd. Mood changes during prednisone bursts in outpatients with asthma. *J Clin Psychopharmacol* 2002;22:55-61.
11. Pies R. Persistent bipolar illness after steroid administration. *Arch Intern Med* 1981;141:1087.

Psychiatric symptoms—especially mania—are a common, unpredictable side effect of corticosteroid use. Taper the corticosteroid, if possible, and add a mood stabilizer or antipsychotic to alleviate symptoms. Consider prophylactic lithium for patients with medical conditions that require frequent or high-dose corticosteroid treatment.

BottomLine

continued



Related resources

- ▶ Brown ES, Chandler PA. Mood and cognitive changes during systemic corticosteroid therapy. *Prim Care Companion J Clin Psychiatry* 2001;3(1):17-21. www.psychiatrist.com/pcc/abstracts/pcc030103.htm.
- ▶ Merrill W. Case 35-1998: use of lithium to prevent corticosteroid-induced mania. *N Engl J Med.* 1999;340:1123.

DRUG BRAND NAMES

- | | |
|--------------------------|----------------------------|
| Amitriptyline • Elavil | Lamotrigine • Lamictal |
| Carbamazepine • Tegretol | Lithium • Eskalith, others |
| Fluoxetine • Prozac | Olanzapine • Zyprexa |
| Gabapentin • Neurontin | Quetiapine • Seroquel |
| Haloperidol • Haldol | Risperidone • Risperdal |

DISCLOSURES

The author reports no financial relationship with any company whose products are mentioned in this article or with manufacturers of competing products.

12. Wada K, Yamada N, Suzuki K, et al. Recurrent cases of corticosteroid-induced mood disorder: a clinical characteristics and treatment. *J Clin Psychiatry* 2000;61:261-7.
13. Narrow WE, Rae DS, Robins LN, Regier DA. Revised prevalence estimates of mental disorders in the United States: using a clinical significance criterion to reconcile 2 surveys' estimates. *Arch Gen Psychiatry* 2002;59:115-23.
14. Brown ES, Chamberlain W, Dhanani N, et al. An open-label trial of olanzapine for corticosteroid-induced mood symptoms. *J Affect Disord* 2004;83:277-81.


15. Blazer DG 2nd, Petrie WM, Wilson WP. Affective psychosis following renal transplant. *Dis Nerv Syst* 1976;37:663-7.
16. Wada K, Yamada N, Yamauchi Y, Kuroda S. Carbamazepine treatment of corticosteroid-induced mood disorder. *J Affect Disord* 2001;65:315-7.
17. Siddiqui Z, Ramaswamy S, Petty F. Quetiapine therapy for corticosteroid-induced mania. *Can J Psychiatry* 2005;50:77-8.
18. Wyszynski AA, Wyszynski B. Treatment of depression with fluoxetine in corticosteroid-dependent central nervous system Sjogren's syndrome. *Psychosomatics* 1993;34:173-7.
19. Brown ES, Frol A, Bobadilla L, et al. Effect of lamotrigine on mood and cognition in patients receiving chronic exogenous corticosteroids. *Psychosomatics* 2003;44:204-8.
20. Terao T, Mizuki T, Ohji T, Abe K. Antidepressant effect of lithium in patients with systemic lupus erythematosus and cerebral infarction, treated with corticosteroid. *Br J Psychiatry* 1994;164:109-11.
21. Terao T, Yoshimura R, Shiratuchi T, Abe K. Effects of lithium on steroid-induced depression. *Biol Psychiatry* 1997;41:1225-6.
22. Ahmad M, Rasul FM. Steroid-induced psychosis treated with haloperidol in a patient with active chronic obstructive pulmonary disease [letter]. *Am J Emerg Med* 1999;17:735.
23. DeSilva CC, Nurse MC, Vokey K. Steroid-induced psychosis treated with risperidone. *Can J Psychiatry* 2002;47:388-9.
24. Falk WE, Mahnke MW, Poskanzer DC. Lithium prophylaxis of corticotropin-induced psychosis. *JAMA* 1979;241:1011-2.
25. Saklad SR. Management of corticosteroid-induced psychosis with lithium. *Clin Pharm* 1987;6:186.
26. Ginsberg DL, Sussman N. Gabapentin as prophylaxis against steroid-induced mania. *Can J Psychiatry* 2001;46:455-6.

Book Byte

Your connection to the latest books for the psychiatric professional

- ! News on book releases
- ! Publishers' reviews
- ! Links to Amazon.com for immediate purchase

Now at www.currentpsychiatry.com



Corticosteroid-induced mania: Prepare for the unpredictable : Current Psychiatry
http://subscribe.imngmedicalmedia.com/onecount/validate_server/getCookie.php?return=http://www.currentpsychiatry.com%2Fhome%2Farticle%2Fcorticosteroid-induced-mania-prepare-for-the-unpredictable%2Fc2e8b70eb48fb07caabb1f0b255060bc.html | via @instapaper.Â Since my #lupus symptoms have eased off for the last year. Iâ€™m struggling with the flares more now. Like Iâ€™ve forgotten how to ignore pain. 0 replies 0 retweets 0 likes.

Corticosteroid-induced mania: Prepare for the unpredictable. Current Psychiatry. June 2006. [www.currentpsychiatry.com/index.php?id=22661&tx_ttnews\[tt_news\]=171412](http://www.currentpsychiatry.com/index.php?id=22661&tx_ttnews[tt_news]=171412).Â Mania triggered by a steroid nasal spray in a patient with stable bipolar disorder. Am J Psychiatry. 1989;146:1076-1077. Corticosteroid-induced side effects are usually managed by tapering off the steroids and treating the psychiatric symptoms.^{2,3} Simply tapering off the steroidsâ€™without additional treatmentsâ€™led to recovery in 33 of 36 patients.² Stopping corticosteroids is not always possible or desirable, however, especially in many medically complicated cases seen by psychiatric consult services. In a recent case, I was asked to see a man, age 69, on the oncology service who was receiving corticosteroids every 2 weeks as part of his chemotherapy. The patient was admitted to the hospital for acute mental stat...Â Mood stabilizers with evidence of benefit in treating corticosteroid-induced mania. Indication. Medication.